



Provisional Maturity Assessment of CrIMSS EDRs

(On Going Efforts with NPP-Aqua Matches and Dedicated RAOBs)



By
Murty Divakarla*, Mike Wilson*, Eric Maddy#,
Xiaozhen Xiong*, Changyi Tan*, Antonia Gambacorta@, and Nick Nalli+
IM Systems Group, Inc. at NOAA/STAR

Xu Liu* and Susan Kizer*
Langley Research Center

Degui Gu*, Xia L Ma*, and Denise Hagan*
Northrop Grumman Aerospace Systems

Chris Barnet*\$ and Mitch Goldberg\$
NOAA/STAR, JPSS

***Core group for CrIMSS-EDR algorithm Implementation, evaluation, Improvements through Discrepancy Reports to JPSS, user support, and, data source for Focus-Day(s) correlative data sets**

STAR in-house Aqua-AIRS retrieval key consultant and data sets

@Initial help on CrIS/ATMS bias-tuning data set

+Coordinator for AEROSE and other ARM/CART dedicated RAOB campaigns

Point of Contact: Chris.Barnet@noaa.gov; Murty.Divakarla@noaa.gov



Presentation Outline




1. Provisional Maturity assessment of CrIMSS EDR Products
 1. Chronology of CrIMSS EDR evaluations from Pre-Launch to Post-Launch and Beta Maturity (Quick Scan)
 2. On going efforts and future plans on validations
 - CrIMSS /Aqua-Matches (Focus Day 05/15/2012)
 - Dedicated RAOBs (Example for the Focus Day 05/15/2012)
2. Evaluation of two different algorithms, the AIRS –Science Team heritage algorithm, and the CrIMSS official EDR Algorithm
 - The attempt here is to present one such example - effect of dust on two different retrieval algorithms – How it was perceived with proxy data, and how it is seen with real observations.
3. Discrepancy Report (DR) Process
 - Algorithm evaluation process, updates, fixes, path way to realization

First-Light CrIMSS 'MW-only' Retrievals


Using SDRs/TDRs, 'the Day-1 Bias Tuning' - Day 11/11/2011

Advantages at NOAA/STAR (from Murty, AMS2012)


MIRS SDR RET




CrIMSS MW-Only (NO Tuning)




CrIMSS-MW Only with MIRS SDR BT




CrIMSS-MW Only with MIRS TDR BT



ECMWF Truth




CrIMSS-MW Only with NUCAPS TDR BT



255 250 261 264 266 269 272 275 278 281 283 286 289 292 295 298

T(p) LAYERS Prof #1 LAT/LON:65S 160W ASC



Thanks to Kevin Garrett for providing MIRS Bias Tuning, and Antonia for providing NUCAPS Bias tuning



Provisional Maturity Assessment

On-going efforts



- » **Evaluate CrIMSS EDR Algorithm Versions**
 - Off-line EDR Algorithm to emulate exact IDPS EDR versions
 - CrIMSS EDRs – Past MX5.3, Present-MX 6.3 (Oct. 2012), and Future (MX7)
 - Updates to CrIMSS EDR algorithm through ADL/G-ADA
 - Evaluate CrIMSS EDRs with truth measurements, identify fixes, suggest EDR algorithm updates through Discrepancy Reports (DRs)
- » **Validations with Focus Day Data Sets (05/15/2012, 09/20/2012) data sets, ECMWF/GFS Fields, and Heritage Algorithm (AIRS) Products.**
 - Matched AIRS/CrIS SDRs and EDRs – A viable path for Provisional Maturity
 - Results Presented are for the Focus Day: 05/15/2012
 - Validations with Dedicated RAOBs once a substantial sample is reached (Example with AEROSPACE RAOB - 05/15/2012)
 - Global RAOB collocations as we have done for AIRS/IASI Validations



Rationale to use AIRS retrievals for CrIMSS Provisional Maturity









- » AIRS Radiances, and AIRS retrievals, especially T(p) , q(p) (and O3) went through Stage-3 validations with a variety of truth data sets. (Murty, JGR-2006, 2008; Dave Tobin, JGR-2006).
- » AIRS V6 (with Pbest ΔT Assimilation QC meets 1K/1Km standard even for cloud-cleared cases. The AIRS defined 'clear' cases (QC = 30hPa down to 750 hPa) RMS difference is pretty close to 0.8K/Km or less (figures follow)
- » Matched Aqua-AIRS retrievals, CrIMSS retrievals and other correlative data sets (ECMWF) can be used to
 - Verify CrIMSS EDR Statistics and Aqua-EDR statistics using ECMWF as the reference (177,000 Matches for the Focus Day 05/15/2012)
 - Globally about 4000 clear cases can be obtained from the AIRS Retrievals, and corresponding matches of Aqua-AIRS/AMSU and CrIS/ATMS SDRs at 3 x3 FOVs, and matched retrievals at FOR resolution can be extracted.
- » This can be further substantiated with validations using 'dedicated RAOBs and/or global RAOB match-up of truth data sets once we achieve a reasonable number of ascents.

500mb Temp (K) Map (Descending)

CLM-2002 NPP-CrIMSS, Aqua-AIRS (V5.9), and ECMWF
(Used in Beta Maturity of CrIMSS)


T(p) RET:25 478.0–535.2 hPa 2012/05/15 DESC CL2002_UPD



222 225 228 231 235 238 241 244 247 250 254 257 260 263 266 269

Temperature (K)

CrIMSS EDRs and Aqua-AIRS Retrievals





Aqua-AIRS and NPP-CrIMSS Versions


» Aqua-AIRS Retrievals

- Version 5.9 uses regression based on PCs trained with ECMWF as the first guess for the final physical retrieval.
- Version 6 uses Neural Network (NN) regression trained with ECMWF as the first guess for the final physical retrieval.
 - Uses pressure dependent QC for each profile, (1) high thresholds for data assimilation applications (pbest), (2) loose thresholds for climate applications (pgood). The yield is higher with pgood, and yield is lower for pbest.
 - We don't have exact QC, but we did somewhat similar
 - » QC = 0 or 1 30 hPa to 750 hPa (higher yield, pgood must be at least 750hPa)
 - » QC = 0 30hPa to 750 hPa (lower yield, pbest must be at least 750 hPa)
 - » Used T(p) QC control for both T(p) and q(p)


» CrIMSS Retrievals.

- Past (MX5.3), Present (MX6.3), Upcoming (MX7)
 - Retrievals used here are some what MX7 equivalent.

AIRS Ret. Versions and RMS Diff (05/15/2012) V5.9, V6(Pgood), V6(Pbest) Global (L+S+C; D+N); Ocean (D+N)



T(p) RMS (K)



Q(p) RMS (%)

Dashed Lines
AIRS V5.9 RET
Dash-Dot Lines
AIRS V6 Pgood
Solid Lines
AIRS V6 Pbest

Red (N= 323000)
GLB (L+S+C, D+N)


V5.9 - Accepted 74%
V6 PGOOD - 86%
V6 PBEST - 32%

Blue (N=193000)
Ocean ALL (D+N)
V5.9 Accepted :75%
V6 PGOOD - 93%
V6 PBEST - 37%
Clear Accepted (3%)
Accepted N= 5,540


Global ALL (L+S+C; D +N) Clear Cases

AIRS Clear Cases - Daytime

Daytime- Land




Daytime - Ocean




AIRS Clear Cases - Nighttime



Nighttime - Land



Nighttime - Ocean



CrIMSS IR+MW vs. ECMWF AIRS V5.9 vs. ECMWF Global Ocean – CLDCLR, Clear, RMS



Solid Lines
CrIMSS IR+MW

Dashed Lines
AIRS V5.9 RET



Blue Ocean
N= 116,000 -
CLDCLR
AIRS:76% dashed
CrIMSS:58% solid

Clear
N= 5,540
AIRS: 3% dashed
CrIMSS solid

T(p) RMS (K)

Q(p) RMS (%)

CrIMSS IR+MW vs. ECMWF AIRS V6 Pgood vs. ECMWF Global Ocean – CLDCLR, Clear, RMS



Solid Lines
CrIMSS IR+MW

Dashed Lines
AIRS **V6** RET
pgood



Blue Ocean
N= 116,000 -
CLDCLR
AIRS: **93%** dashed
CrIMSS: **58%** solid

Clear
N= 5,520
AIRS: **3%** dashed
CrIMSS solid

T(p) RMS (K)

q(p) RMS (%)

CrIMSS IR+MW vs. ECMWF AIRS V6 Pbest vs. ECMWF Global Ocean – CLDCLR, Clear, RMS



Solid Lines
CrIMSS IR+MW

Dashed Lines
AIRS V6 RET
pbest



Blue Ocean
N= 116,000 -
CLDCLR
AIRS:43% dashed
CrIMSS:58% solid

Clear
N= 5,538
AIRS: 3% dashed
CrIMSS solid

T(p) RMS (K)

q(p) RMS (%)

CrIMSS IR+MW vs. ECMWF AIRS V6 pbest vs. ECMWF Global (D+N) : (L+S+C); CLDCLR, Clear RMS



Solid Lines
CrIMSS IR+MW

Dashed Lines
AIRS V6 RET
pbest

Global ALL
N=177,000
AIRS: 38% dashed
CrIMSS: 53% solid
CLDCLR

Clear
N: 5,277
AIRS: (3%) dashed
CrIMSS: solid
5331

T(p) RMS (K)



q(p) RMS (%)



Global Ocean - ± 60 LAT, Night Time

AIRS V6(pbest), CrIMSS CLDCLR, Clear, STDEV

Clear: N:3,308 'Dedicated' Matches (pg.



Solid Lines
CrIMSS IR+MW

Dashed Lines
AIRS V6 RET
pbest

Blue Ocean
N= 52,000 -
CLDCLR
AIRS:51% dashed
CrIMSS:59% solid
with
AIRS WGT
Dash-dot

Clear
N= 3,308
AIRS: 3% dashed
CrIMSS solid

T(p) RMS (K)

Q(p) RMS (%)

- » EDR Evaluations with ECMWF
 - First Stage CrIMSS-MW RET (ATMS) vs. AMSU-MIT Ret
 - ATMS Retrievals are better than AMSU Retrievals (Plots not shown)
 - IR+MW Stage CrIMSS RET vs. AIRS-PR
 - Clear (6326 ; 5520 ocean; 3308 ocean, night) and CLDCLR cases
 - » Very encouraging results for matched clear/cloud-cleared cases
 - » EDR Evaluations of clear cases are similar to AIRS
 - » CrIMSS QC flags – A revisit and a DR in the process.
 - Retrieval ability over dust-free/dusty areas
- » This Matched AIRS-CrIS data set is very useful and can lead us to provisional maturity very easily and can be considered as ‘Dedicated Matches’ synonymous to Dedicated RAOBs that are very sparse.
- » Improve CrIS/ATMS empirical bias tuning with clear cases as defined by AIRS algorithm and AIRS/MODIS clear matches


Aerospace RAOB (Kauai, Hawaii) Location Specifics with ATMS BTs

ATMS - Sub-Pixel FOV Variability with in FOR

Example Region




ATMS (CH#1) Field of Regard BT Difference





Field of Regard (Sub-Pixel Variability)

ATMS-TDRs




CH #1 BT Difference (FOR) (obs-cal)




Aerospace RAOB (Kauai, Hawaii) Location Specifics with ATMS BTs CH#3

IRS/CrIMSS Ret Matches
Kauai Hawaii RAOB Location on ATMS Brightness Temperature CH # 3
IRS Acc. - Nbr CLR(Red Dots):113, Nbr CLDCLR(Black Dots):877




Aerospace RAOB (Kauai, Hawaii) Location Specifics Total PCW CrIMSS




Aerospace RAOB (Kauai, Hawaii) Location Specifics Total PCW ECMWF

AIRS/CrIMSS Ret Matches
Kauai Hawaii RAOB Loc. Total PCW(mm) from ECMWF
AIRS Acc. - Nbr CLR(Red Dots):113, Nbr CLDCLR(Black Dots):877




Aerospace RAOB (Kauai, 22.05N, 159.78W, Hawaii) Location Specifics with ATMS BTs CH#1


AIRS-V5.9/CrIMSS Ret Matches - Kauai, Hawai RAOB with ATMS BT CH # 1
AIRS Acc. CLR(Red Dots) CLDCLR(Black Dots):877
Subset CLDCLR/CLEAR Cases with Distance to AERO-RAOB & With QC



Aerospace RAOB (Kauai, 22.05N, 159.78W, Hawaii) Location Specifics with ATMS BTs CH#1



CrIMSS Ozone and OMPS Ozone (Provided by Jianguo Niu, Larry Flynn, STAR)




October 2, 2012 (Top); October 16, 2012 (Bottom)
Towards TOAST of CrIMSS and OMPS Ozone
(TOAST: Total Ozone Analysis from SBUV and TOVS)

Aqua-AIRS Ozone Validation, (Murty - JGR, 2008)

WOUDC O3(p), TO3 BD Measurements

Synergetic Use of A-Train Satellite Data Sets (Aura-OMI, N16-SBUV)






Matched WOUDC BD Measurements Used in Validating AIRS Total Ozone N~4096 Bias (4%, OMI (3%), RMS (8%)

AIRS TO3 vs. BD
Global
8% RMS
4% Bias



AIRS Total Ozone
February 2005

Annual Cycle
AIRS - 2005




OMI Total Ozone
February 2005


Annual Cycle
OMI - 2005




Divakarla et. al., 'Evaluation of AIRS Ozone Profiles and Total Ozone Retrievals with Matched O3SNDs, ECMWF and OMI Retrievals, J. Geophys. Res., 113, D15308, doi:10.1029/2007JD009317. <http://www.agu.org/pubs/crossref/20082007JD009317.shtml>



AIRS
1800 Km
Swath



OMI
2600 Km
Swath




SBUV
Nadir



Summary

- » We are in the process of optimizing MX-7 CrIMSS EDR Algorithm Emulation and going through a thorough evaluation/implementation of many Discrepancy Reports. Results shown here will be improved substantially with all the anticipated changes.
- » This AIRS/CrIS matches provide a simple pathway for CrIMSS EDR Provisional Maturity . The AIRS-CrIMSS SDR/EDR Matches (Ocean, Night Time, ± 60 Lat, N: 3,300 for 05/15/2012) can be considered as ‘Dedicated Aqua Matches’ synonymous to ‘Dedicated RAOBs’ . Extending this to other Focus Days could provide Set of ‘Golden Focus Day Data Sets’ to
 - Validate CrIMSS vis-à-vis AIRS Retrievals with Matched ECMWF/ Dedicated RAOBs
 - Refine and enhance CrIS/ATMS empirical bias tuning
 - Provide the best possible evaluation of two different algorithms, the AIRS –Science Team heritage algorithm, and the CrIMSS official EDR Algorithm) - Slides Follow →
- » The attempt here is to present one such example - effect of dust on two different retrieval algorithms – How it was perceived with proxy data, and how it is seen with real observations.

On-Going STAR Activities In Collaboration with Cal/Val Team Members



As Directed by:
Chris Barnet, CrIMSS Cal/Val Team Lead

Topic:
**Discrepancy Reports Generation
Coordinated Efforts to Expedite
Algorithm Change Process**

Degui Gu, Xia L Ma, and Denise Hagan
Northrop Grumman Aerospace Systems



Xu Liu and Susan Kizer
Langley Research Center

*Mike Wilson, Murty Divakarla, Bigyani
das, Changyi Tan, and Xiaozhen Xiong,*
STAR, NOAA/NESDIS

Wael Ibrahim
Raytheon
Other Cal/Val Teams
Other Teams (Richard Cember)

How Do We Move forward

- “what and how” of proposed changes can be expedited through Algorithm Change Process
- Make sure the DR submitted is worthy of consideration
 - Scientific evaluation of the DR.
 - Provide preemptive answers to the queries expected in the implementation process



DR Table: MX6.5 to MX7.0

DR #	Files Affected	LUT Affected	Description
4926	calcCrimssProfiles.f set_irmw_invert.f		This change fixes handling of forward model errors. The current code has errors in computing total error in radiance residuals.
4942	fovsel.f		The current code identifies scenes as clear when they are actually partly cloudy. This change causes tighter scene detection for clear scenes.
4943		CrIMSS-IR-NOISE CrIMSS-IR-ATM-NOISE	This change upgrades the sensor and forward model LUTs for the CrIS instrument. The current LUTs are pre-launch values.
4945	calcCrimssProfiles.f		This change allows the difference between skin temperature and surface temperature to be larger for daytime land.
4946	setCovBack.f		This change optimizes the climatology LUT. The criteria for choosing warm ocean is loosened through this change, resulting in the warm ocean climatology being chosen more frequently.
4958		CrIMSS-EDR-AC	This change optimizes the QC flags. Currently, the QC flag for the microwave portion of the combined run is too strict. This change increases the threshold for this flag from 2 to 4, making values pass more frequently.



What We have Presented.

1. Provisional Maturity assessment of CrIMSS EDR Products
 1. Chronology of CrIMSS EDR evaluations from Pre-Launch to Post-Launch and Beta Maturity (Quick Scan)
 2. On going efforts and future plans on validations
 - CrIMSS /Aqua-Matches (Focus Day 05/15/2012)
 - Dedicated RAOBs (Example for the Focus Day 05/15/2012)
2. Evaluation of two different algorithms, the AIRS –Science Team heritage algorithm, and the CrIMSS official EDR Algorithm
 - The attempt here is to present one such example - effect of dust on two different retrieval algorithms – How it was perceived with proxy data, and how it is seen with real observations.
3. Discrepancy Report (DR) Process
 - Algorithm evaluation process, updates, fixes, path way to realization
 - **Thank You.**



End of Presentation

